




The AGN Physics NuSTAR program

Giorgio Matt


(Universita' Roma Tre, Italy)

On behalf of the NuSTAR AGN Physics WG

NuSTAR is the **first** focusing hard X-ray satellite




Coded Aperture Optics:
high background, large detector



Grazing Incidence Optics:
low background, compact detector

Collecting Area



*NuSTAR two-telescope total
collecting area*

Sensitivity comparison

INTEGRAL (ISGRI) ***~0.5 mCrab***
(20-100 keV)
with >Ms

Swift (BAT) ***~0.8 mCrab***
(15-150 keV)
with >Ms

NuSTAR ***~0.8 μCrab***
(10-40 keV)
In 1 Ms

1 Ms Sensitivity

3.2×10^{-15} erg/cm²/s

(6 – 10 keV)

1.4×10^{-14} (10 – 30 keV)

Timing

relative 100 microsec

absolute 3 msec

Imaging

HPD 58"

FWHM 18"

Localization 2" (1- σ)

Spectral response

energy range 3-79 keV

threshold 2.0 keV

ΔE @ 6 keV 0.4 keV FWHM

ΔE @ 60 keV 1.0 keV FWHM

Field of View

FWZI 12.5' x 12.5'

FWHI 10' @ 10 keV

8' @ 40 keV

6' @ 68 keV

Target of


Opportunity

response <24 hr (reqmt)

typical 6-8 hours

80% sky accessibility

High-Energy Missions in Orbit: comparison of pixel scales



Imaging

Cas A supernova remnant

INTEGRAL ISGRI



$E > 15$ keV

NuSTAR Image

Red : 4.5 – 5.5 keV


Green: 8 – 10 keV

Blue: 10 – 25 keV




Grefenstette et al. (2014)

Imaging



X-Ray Telescopes & the Electromagnetic Spectrum



Baseline Science Mission

- Small Explorer Mission (SMEX), launched in June 2012
- As typical for an Explorer, all baseline observations led by the science team
- After the current initial calibration period has been completed, observations became public through HEASARC two months after a data set is completed
- 1.5 Ms of NuSTAR made available for coordinated observations with XMM in AO13 (with a factor 6 oversubscription)
- Mission extended to 2015-16 with allocated budget for two more years
- GO program will start in April 2015 (deadline for proposals: November 25, 2014)
- ~140-person international science team broken into 13 science working groups:

Science Working Groups

Science Group




Galactic Survey
Supernovae and ToOs
Supernova Remnants and PWN
Magnetars and RPP
Galactic Binaries
Ultraluminous X-ray sources
Extragalactic Surveys
Blazars
Obscured AGN
AGN Physics
Galaxy Clusters
Starburst Galaxies
Solar Physics

Working Group Chair

Chuck Hailey
Steve Boggs
Fiona Harrison
Vicky Kaspi
John Tomsick
Fiona Harrison
Daniel Stern
Greg Madejski/Paolo Giommi
Daniel Stern
Giorgio Matt
Allan Hornstrup/Silvano Molendi
Ann Hornschemeier
David Smith

AGN Physics: Scientific rationale

- ❖ Determine the physical parameters of the hot corona (temperature, optical depth)
- ❖ Measure the spin of the Black Hole
- ❖ Search for similarities and differences between radio quiet and radio loud AGN



RQ objects in the AGN Physics NuSTAR Program

- 4 sources observed simultaneously with XMM [**Swift J2127.4+5654, MCG-6-30-15, 3C120, Ark 120**] for BH spin and corona T. **Ark 120** re-observed recently in coordination with XMM and Chandra.
- 2 sources observed simultaneously with Suzaku [**IC4329A, NGC4151**] for BH spin and corona T
- **MCG-5-23-16** observed twice, the second time simultaneously with Suzaku
- **Mrk 335** observed, in coordination with Suzaku, during an extended low state
- A monitoring campaign studied the spectral variability of **NGC4051**
- NuSTAR joined the XMM-led monitoring campaign on **NGC5548** and on **PDS 456**
- **1H0707+495** and **Fairall 9** observed recently (the latter simultaneously with XMM)

Content of this talk



- The coronal parameters in **IC4329A**, **MCG-5-23-16** and **SwiftJ2127.4+5654**
- The relativistic reflection in **NGC1365** (*Risaliti et al., 2013, Walton et al. 2014*) and the BH spin of **SwiftJ2127.4+5654** and **MCG-6-30-15** (*Marinucci et al., 2014a,b*)
- The hard X-rays time lags of **MCG-5-23-16** (*Zoghbi et al., 2014*)
- The extreme relativistic reflection in **Mrk 335** (*Parker et al. 2014*)
- The soft excess of **Ark 120** (*Matt et al., 2014*)

Coronal parameters

Primary hard X-ray emission likely due to Comptonization in a hot corona → quasi-exponential high energy cutoffs expected

Evidence for high energy cutoffs in BeppoSAX and XMM - INTEGRAL samples

NuSTAR is providing for the first time source-dominated obs above 10 keV → coronal parameters




Coronal parameters


Primary hard X-ray emission due to Comptonization in a hot corona → high energy cutoffs expected

Evidence for high energy cutoffs in BeppoSAX and XMM - INTEGRAL samples

NuSTAR is providing for the first time source-dominated obs above 10 keV → coronal parameters



(Perola et al. 2002)




(Malizia et al. 2014)

Coronal parameters

Primary hard X-ray emission due to Comptonization in a hot corona → high energy cutoffs expected


Evidence for high energy cutoffs in BeppoSAX and XMM - INTEGRAL samples

NuSTAR is providing for the first time source-dominated obs above 10 keV → coronal parameters




Swift J2127.4+5654 (Marinucci et al. 2014)

**$kT \sim 68/53$ keV $\tau \sim 0.35/1.35$
(slab/sphere)**



**IC4329A
(Brenneman et al. 2014)**



**$kT \sim 61/50$ keV $\tau \sim 0.7/2.35$
(slab/sphere)**



**Ark 120
(Matt et al. 2014)**

Coronal parameters



The best case so far: MCG-5-23-16 (Balokovic et al., submitted)




The relativistic reflection in NGC1365



Prograde Rotation Model




Foreground Obscuration Model



The relativistic reflection in NGC1365

Observed simultaneously by XMM and NuSTAR.



Both absorption and reflection models fit well the XMM data, but only reflection also the NuSTAR data (Risaliti et al. 2013)



The relativistic reflection in NGC1365

Observed simultaneously by XMM and NuSTAR.



Consistent results are found in all observations, despite huge differences in the absorption parameters (Walton et al. 2014)



The relativistic reflection in NGC1365


Observed simultaneously by XMM and NuSTAR.

Consistent results are found in all observations, despite huge differences in the absorption parameters (Walton et al. 2014)



BH spin measurements

The broad band provided by NuSTAR + XMM (or Suzaku) allows a good estimated of the continuum spectrum, and so a robust measurements of the BH spin via relativistic effects on the iron line and the reflection component




Spin ~ 1 confirmed in MCG-6-30-15 (Marinucci et al. 2014b)

SwiftJ2127.4+5654
(Marinucci et al. 2014a)


The hard X-ray time lag in MCG-5-23-16

Soft time lags observed in many AGN (e.g. Fabian et al. 2009, De Marco et al. 2013, Uttley et al. 2014) --> Reflection from inner disc
More recently, reverberation of iron lines have also been observed (e.g. Zoghbi et al. 2012, Kara et al. 2014)

Compton hump reverberation expected !!





XMM (Zoghbi et al. 2013)





NuSTAR (Zoghbi et al. 2014)

Mrk 335: Relativistic effects within 2 R_g from the event horizon?




The source was found in a very low flux state (Parker et al. 2014).

Mrk 335: Relativistic effects within 2 R_g from the event horizon?



The spectrum is well fitted by an almost pure relativistic reflection component. Applying a lamp-post geometry, a very small height is found, as well as a high BH spin (Parker et al. 2014)

The soft excess of Ark 120




Bright, “bare” Seyfert 1 galaxy

Fit with NuSTAR data only (power law + reflection + iron line)

No High Energy Cutoff detected

Extrapolation to XMM shows strong excess




(Matt et al. 2014)



The soft excess of Ark 120



**XMM: no obvious evidence for rel. Line
(differently from a previous Suzaku obs,
Nardini et al. 2011)**

**Soft excess with a simple power law or with
a Comptonization model give comparable
fits to the XMM spectrum, but
very different extrapolation to NuSTAR
(cold and ionized reflection included in the fit)**



The soft excess of Ark 120

Indeed, the broad-band best fit is with a Comptonization model for the soft excess. A *cutoff p.l.*, *compTT*, *nthcomp* or *optxagnf* provide fits of comparable quality.



Optxagnf (Done et al. 2012) is a disk/corona emission model which assumes a thermal disk emission outside the coronal radius, and soft and hard Comptonization inside.

Extrapolating the best fit X-ray model to the OM UV data, an estimate of the black hole spin is possible

Summary and Outlook

NuSTAR is providing AGN spectra of unprecedented quality above 10 keV

A joint XMM-NuSTAR observing program also in AO14

*A **GO program** will start in april 2015
(proposals due by November 25, 2014)*